

Competing risks in epidemiology: Possibilities and pitfalls

Per Kragh Andersen
Ronald B. Geskus
Hein Putter

Research Report 11/2

Department of Biostatistics
University of Copenhagen

Competing risks in epidemiology: Possibilities and pitfalls

Per Kragh Andersen

University of Copenhagen

Denmark

Ronald B. Geskus

Academic Medical Center and Public Health Service Amsterdam

The Netherlands

Hein Putter

Leiden University Medical Center

The Netherlands

May 11, 2011

Abstract

In studies of all-cause mortality the fundamental epidemiological concepts of rate and risk are connected through a well-defined one-to-one relation. An important consequence of this relation is that regression models such as the proportional hazards model that are defined

through the hazard (the rate) immediately dictate how the covariates relate to the survival function (the risk). This introductory paper reviews the concepts of rate and risk and their one-to-one relation in all-cause mortality studies and introduces the analogous concepts of rate and risk in the context of competing risks, the cause-specific hazard and the cause-specific cumulative incidence function. The key feature of competing risks is that the one-to-one correspondence between cause-specific hazard and cumulative incidence, between rate and risk, is lost. This fact has two important implications. First, the naive Kaplan-Meier which takes the competing events as censored observations, is biased. Second, the way in which covariates are associated with the cause-specific hazards may not coincide with the way these covariates are associated with the cumulative incidence. An example with relapse and non-relapse mortality as competing risks in a bone marrow study is used for illustration.

Key words: censored data, competing risks, regression models, survival analysis.

1 Introduction

Epidemiology deals with the occurrence of diseases in populations when observed over time, and the frequency with which diseased cases occur are measured using the concepts of *risk* and *rate*. Standard text books in epidemiology (e.g. Rothman¹⁰, Chapter 3, dos Santos Silva⁴, Section 4.2) typically define the risk as the fraction $\frac{D}{N}$ of N originally disease-free individuals in the population who develop the disease over a specified follow-up period, say, from time 0 to time t . Note that the risk must necessarily increase with

t . On the other hand, the rate would typically be defined as the number, D , of individuals in the population who develop the disease during a specified follow-up period (from 0 to t) divided by the amount of person-time, Y at risk observed when following disease-free individuals from the population from 0 to t . The rate $\frac{D}{Y}$ may increase, stay roughly constant, or decrease when varying the length, t of the follow-up period.

The statistical counterpart of a risk is a *probability*. Thus, if $F(t)$ denotes the probability that a randomly selected disease-free individual gets the disease before time t then the risk $\frac{D}{N}$ estimates $F(t)$ if all N disease-free individuals in the population are followed from 0 to t . However, in most follow-up studies there will inevitably be loss to follow-up, *censoring*, and $F(t)$ must then be estimated using more complicated techniques able to account for censoring.

The statistical discipline that deals with censored follow-up data is *survival analysis* and in the next paragraphs we will summarize basic (perhaps well-known) features of survival analysis. We will do that in the context where the event under study (“the disease”) is all-cause mortality, that is, an event which will occur with probability one if the follow-up period is sufficiently long (t is “large”). However, our motivation for doing this is to set the scene for the situation where observation of the disease under study may be preceded by other events, the occurrence of which prevents us from observing the disease of interest. This *competing risks* situation (which is the rule rather than the exception in epidemiological follow-up studies) is the topic for the present paper. We shall discuss which concepts from classical survival analysis (i.e., studies of all-cause mortality) immediately extend to

competing risks and we shall discuss when to be more careful.

2 Survival analysis

In survival analysis the object is the time elapsed from an initiating event, e.g. the onset of some disease, to death. If time to death was observed for every one in the sample then the probability $F(t)$ of dying before time t (the *cumulative incidence function*), as explained above, could be estimated as the relative frequency of survival times less than t . However, the challenge is to estimate $F(t)$ based on incomplete data, i.e. to make inference on the underlying, potentially completely observed population in the presence of censored observations. For this to be feasible, censoring must be *independent*, that is, an individual censored at time t should be representative for those still at risk at that time. In other words, those censored should *not* be individuals with systematically high or low risk of dying. Under independent censoring, $F(t)$ may be estimated by $1 - \hat{S}(t)$ where $\hat{S}(t)$ is the *Kaplan-Meier estimator* for the probability $S(t) = 1 - F(t)$ of surviving time t . The Kaplan-Meier estimator at time t is a product with a factor for each failure time before t . The factor at failure time s is $(1 - \frac{D_s}{N_s})$ where D_s is the number of failures observed at s (often $D_s = 1$), and N_s the number of individuals in the study still *at risk*, i.e. alive and uncensored, at time s .

The concept in survival analysis that corresponds to the *rate* is the *hazard function* $h(t)$. This has the interpretation that for a small interval from time t to time $t + d$, $h(t) \cdot d$ is approximately the conditional probability of death before time $t + d$ given survival until time t . Thus, the hazard function provides a dynamical (“local in time”) description of how the instantaneous

risk of failing varies. The epidemiological rate $\frac{D}{Y}$ mentioned above is then a sensible estimate for the hazard function if this is roughly time-constant, i.e. when $h(t) = h$ then $\hat{h} = \frac{D}{Y}$ estimates h .

In survival analysis there is a simple “one-to-one” correspondence between the hazard function and the survival function, derived in Box 1. For any

If $h(t)$ is constant then this is seen as follows. (If $h(t)$ is not constant then a similar argument applies.) We divide the interval from 0 to t into K small intervals each of length d , that is $t = Kd$. Now, in order to survive time t one must first survive the first little interval (from 0 to d). This happens, by definition of the hazard, with probability $1 - h \cdot d$. Given survival through the first little interval, the probability of surviving the next is also $1 - h \cdot d$, that is, the probability of surviving the first two intervals is $(1 - h \cdot d)(1 - h \cdot d) = (1 - h \cdot d)^2$. Continuing this argument it is seen that the probability, $S(t)$, of surviving time $t = Kd$, that is, of surviving K little intervals, is $(1 - h \cdot d)^K = (1 - h \cdot \frac{t}{K})^K$. When K is large (and, thereby, d is small) this expression equals e^{-ht} , that is, the negative exponential of the *cumulative hazard function* ht .

Box 1: Relation between hazard and survival function in survival analysis

hazard function (not necessarily constant) the relationship is

$$S(t) = e^{-H(t)} \text{ or } F(t) = 1 - e^{-H(t)} ,$$

where $H(t)$ is the cumulative hazard function at time t . It follows that, for given hazard function $h(t)$ one may compute the survival function $S(t)$ (or the cumulative incidence function $F(t)$), and vice versa: for given cumulative incidence the hazard may be computed. Under independent censoring the cumulative hazard at time t may be estimated by the *Nelson-Aalen estimator*.

This is a sum with a term for each failure time before t , the term at failure time s being $\frac{D_s}{N_s}$. Note how the “one-to-one correspondence” between rate and risk is reflected in the Kaplan-Meier and Nelson-Aalen estimators which are both based on the same basic pieces of information: the number of failures, D_s and the number at risk, N_s at each failure time, s .

This has important consequences for the analysis of survival data because models for the hazard function, e.g., the Cox regression model² which is very frequently used, immediately imply models for the cumulative incidence $F(t)$. Thus, if a Cox regression model is fitted for the hazard function and if, based on this Cox model, presence of a certain factor is seen to be associated with a higher hazard function then presence of the factor is also associated with a higher cumulative incidence. The interpretation of the parameter estimated in a Cox regression model is a *hazard ratio*.

Example

For illustration we use data of the European Group for Blood and Marrow Transplantation (EBMT). The data consists of all chronic myeloid leukemia (CML) patients, having received an allogeneic stem cell transplantation from an HLA-identical sibling or a matched unrelated donor during the years 1997–2000. Patients had to be Philadelphia positive, transplanted with bone marrow or peripheral blood, and 18 years of age or older, leaving 3982 patients. Median follow-up was 8.5 years. An important and very predictive risk score is the EBMT risk score by Gratwohl et al.⁸, originally taking values 0 through 7, and often (also here) for convenience grouped into five distinct groups, with EBMT risk score 0, 1 ($n = 506$), 2 ($n = 1159$), 3 ($n = 1218$), 4

($n = 745$), and 5, 6, 7 ($n = 354$). Failure from transplantation may either be due to relapse, or to non-relapse mortality (NRM). Often these two endpoints are taken together to define what is called relapse-free survival (RFS), which is the time from transplantation to either relapse or death, whichever comes first. Table 1 shows counts and observed percentages of these events in each of the EBMT risk groups. The censored patients were alive without relapse at the end of their follow-up.

[Table 1 about here.]

Figure 1 shows both the Nelson-Aalen estimates of the cumulative hazards (left) and the Kaplan-Meier estimates of the survival curves (right) for RFS for each of the five risk groups.

[Figure 1 about here.]

Cox regression for relapse-free survival gives hazard ratios (95% confidence intervals) of 1.27 (1.08–1.48), 1.61 (1.38–1.88), 2.08 (1.77–2.45), and 3.26 (2.73–3.91) of EBMT risk groups 2, 3, 4, and 5/6/7, respectively, with respect to the reference risk group of 0/1. Clearly, higher EBMT risk scores imply higher rates of the composite endpoint relapse or death, consistent with the left panel of Figure 1. Note that, due to the one-to-one correspondence between rate and risk, higher EBMT score also implies higher risk of relapse or death, i.e. lower relapse-free survival curves.

In conclusion, survival data (all-cause mortality data) may be characterized either by the “global parameter”, the cumulative incidence function $F(t)$ or by the “local parameter”, the hazard function $h(t)$. These two ways of characterization are equivalent due to their one-to-one correspondence.

3 Competing risks

Suppose now that the event of interest is the onset of a given disease but that, obviously, individuals may die without getting the disease. We may then be interested in the risk or probability of getting the disease in a given follow-up period from 0 to t or in the rate or hazard of getting the disease. A naive analysis inspired by the methods for survival analysis outlined above could consider death without the disease as “independent censoring”, thereby aiming at making inference for an underlying, potentially completely observed population. However, that population would be one without “censoring”, that is, a purely hypothetical population where individuals could not die without the disease. A much more satisfactory approach, to be outlined in the following, is one where one acknowledges that individuals may die without the disease and where inference for disease risks and rates are made *in the presence of the competing risk of dying*.

Define $F_1(t)$ as the probability (cumulative incidence) of getting the disease before time t . Define, further, the *cause-specific hazard function* for the disease, $h_1(t)$ as follows: $h_1(t) \cdot d$ is (approximately, when d is small) the conditional probability of getting the disease before time $t + d$ given that the individual is alive and disease-free up to time t . Now, in each little interval from t to $t + d$ between time 0 and time t where the individual is still at risk (alive and disease-free) he or she both has the possibility of getting the disease or dying (without having got it). Therefore, the cumulative incidence of getting the disease not only depends on $h_1(t)$ but also on the hazard of dying. This *cause-specific hazard of death*, $h_2(t)$ is defined similarly to $h_1(t)$; $h_2(t) \cdot d$ is (approximately, when d is small) the conditional probability of

dying before time $t + d$ given that the individual is alive and disease-free up to time t . The consequence is that in the presence of the competing risk of death there is no longer a one-to-one correspondence between the (cause-specific) hazard (the rate $h_1(t)$) and the probability (cumulative incidence) (the risk $F_1(t)$) for the disease and in order to compute the cumulative incidence, the cause-specific hazard for the competing event is also needed. The relationship, derived in Box 2, is

$$F_1(t) = \int_0^t e^{-H_1(s)-H_2(s)} \cdot h_1(s) \cdot ds .$$

This expression shows that, via the factor involving $H_2(t)$, the cumulative incidence for one failure cause (here: the disease) depends on the rate (cause-specific hazard) for the competing cause (here: death without the disease). This is the key feature of competing risks. There is no longer a one-to-one correspondence between cumulative incidence and cause-specific hazard (“rate and risk”). This fact has two important implications:

1. a naive estimator for the cumulative incidence $F_1(t)$ which only studies cause 1 events (disease cases), e.g. 1 minus the Kaplan-Meier estimator based only on disease events and treating deaths as independent censorings is (upwards) *biased*;
2. the way in which the cumulative incidence $F_1(t)$ is associated with covariates may not coincide with the way in which the cause-specific hazard $h_1(t)$ is associated with covariates, but will also depend on the association between covariates and the cause-specific hazard for the competing event $h_2(t)$.

Divide the interval from 0 to t into many small intervals each of length d . To get the disease before time t it must occur in exactly one of these small intervals and the probability of getting the disease before time t , i.e. the cumulative incidence $F_1(t)$, is therefore the sum of the probabilities of getting the disease exactly in each little interval. The probability of getting the disease in the little interval from time s to time $s + d$ is the probability of being alive and disease-free until time s times the conditional probability of getting the disease between s and $s + d$ given alive and disease-free at s . The latter conditional probability is, by the definition of the cause-specific hazard, (approximately) equal to $h_1(s) \cdot d$ while the probability of being alive and disease-free (that is, the probability of staying event-free) until time s is (by an argument similar to the one given above for survival data) equal to $e^{-H_1(s)-H_2(s)}$. Here $H_1(s)$ and $H_2(s)$ are the cumulative cause-specific hazards for the two competing events, disease and death without the disease. The result is that $F_1(t)$ is a sum of terms given by $e^{-H_1(s)-H_2(s)} \cdot h_1(s) \cdot d$ where the sum is over all small intervals between 0 and t . Mathematically, this sum is the integral

$$F_1(t) = \int_0^t e^{-H_1(s)-H_2(s)} \cdot h_1(s) \cdot ds.$$

Box 2: Relation between cause-specific hazards and cumulative incidences in competing risks

Example, continued

To illustrate the first point, consider the highest risk group, with EBMT risk score 5, 6, 7. Figure 2 shows the naive Kaplan-Meier estimates for relapse (censoring patients that died before relapse) and for non-relapse mortality

(censoring patients with relapse) for this highest risk group. The estimate of NRM is shown as a survival curve (starting at 1 and decreasing), the estimate of relapse as an incidence curve (starting at 0 and increasing).

[Figure 2 about here.]

The estimated five-year probabilities of relapse and NRM, obtained from these naive Kaplan-Meiers, are 0.515 and 0.569, respectively. It is clear that these can never be unbiased estimates of the probabilities of relapse and NRM at five years, since they add up to more than 1. This is impossible, since relapse and NRM are mutually exclusive events. The correct estimates, using the relation derived in Box 2, are shown in Figure 3. The previously obtained naive Kaplan-Meier estimates are shown in grey.

[Figure 3 about here.]

The estimated five-year probabilities of relapse and NRM, obtained from this relation, are 0.316 and 0.475, respectively, and the five-year RFS probability is $1 - 0.316 - 0.475 = 0.209$, exactly as obtained in the previous section (see also Figure 1).

Next we turn to the second point: the way covariates affect hazards may be different from the way they affect cumulative incidences. Figure 4 shows Nelson-Aalen estimates of the cumulative cause-specific hazards of relapse and non-relapse mortality for each of the five EBMT risk groups.

[Figure 4 about here.]

The overall picture is that higher EBMT risk score implies higher cause-specific hazards. This is particularly clear for non-relapse mortality; the same

is true in general for relapse, but the two lowest risk groups, those with risk scores 0, 1, and with risk score 2, are approximately equal. Table 2 shows the hazard ratios and 95% confidence intervals of the EBMT risk groups for relapse and non-relapse mortality, obtained from two Cox proportional hazards models, one for relapse (censoring patients dying without relapse), the other for non-relapse mortality (censoring patients with relapse).

[Table 2 about here.]

Although, based on Figure 4, one could question the validity of the proportional hazards assumption, we see from Table 2 that the cumulative cause-specific hazards for relapse are quite similar for risk scores 0, 1 and for risk score 2. If anything, the risk group with score=2 has slightly higher rate. Note that while, as argued above, the Kaplan-Meier estimator should not be used for *risk estimation* in the presence of competing risks, we have used both the Nelson-Aalen estimator and the Cox regression model for the cause-specific *rates*. We will return to an explanation of this apparent paradox in the next section.

Based on these Cox models for the cause-specific hazards for relapse and NRM, we calculated, again using the relation derived in Box 2, the model-based cumulative incidences for relapse and NRM for each of the EBMT risk groups. The results are shown in Figure 5.

[Figure 5 about here.]

Comparing these cumulative incidences of relapse for the two lowest risk groups, we notice a striking thing: the cumulative incidence of relapse is *lower* for the group with EBMT risk score 2, compared to the group with

EBMT risk score 0, 1. Contrast this with what we saw earlier, namely that there is no difference on the cause-specific hazard of relapse between these two lowest risk groups (if anything, the rate for EBMT risk score 2 is higher). The example thus shows that the effect of EBMT risk score on the risk of relapse, the cumulative incidence, is different from its effect on the rate, the cause-specific hazard. The fact that the cumulative incidence of relapse is lower for the EBMT risk score 2 group, compared to the EBMT risk score 0, 1 group, even though the cause-specific hazard of relapse is (somewhat) higher, can be seen as follows. Ignoring true censorings for the moment, the *rate*, the cause-specific hazard of relapse, acts on those individuals still at risk, i.e. on those alive without relapse. The Cox model tells us that, at any point in time, this rate is higher for the EBMT risk score 2 group. But the cause-specific hazard of the competing event, non-relapse mortality, is also higher for the EBMT risk score 2 group, compared to the EBMT risk score 0, 1 group, and the difference here is much larger. That means that over time, the risk set of the EBMT risk score 2 group decreases much more quickly than that of the EBMT risk score 0, 1 group. As a result, even though, relative to the size of the risk set, more individuals will have a relapse in the EBMT risk score 2 group, in absolute size there will in fact be fewer individuals with a relapse. Hence, the *risk*, the cumulative incidence of relapse, will be lower in the EBMT risk score 2 group.

4 Risk and rate models for competing risks

Recall the useful interpretation of the hazard rate in all-cause mortality studies from Section 2: $h(t)$ is approximately the *instantaneous risk per time unit*

of failure at time t given survival till just before t . Thereby, the parameters in the Cox regression model (see Example in that section) are hazard rate ratios. The interpretation carries over verbatim to the cause-specific hazard rate as introduced in Section 3: $h_1(t)$ is approximately the *instantaneous risk per time unit* of failure at time t *from cause 1* given survival till just before t . Similarly, the parameters in Table 2 are ratios between cause-specific hazards. Given the close similarity in interpretation, it is perhaps not entirely surprising that estimation of hazard rate parameters carries over to estimation of parameters in models for cause-specific hazards and, indeed, both the Nelson-Aalen estimator and the Cox regression model may be applied for cause-specific hazards in a fashion completely analogously to studies of all-cause mortality by censoring individuals failing from competing causes. We used this fact for the analyses in the Example in Section 3. The intuitive explanation is that both types of hazard rates describe what happens *locally in time* among individuals still at risk. The formal explanation, given in Box 3, is that the likelihood factorizes.

To estimate risks, the results from a rate model may be plugged into the relationship derived in Box 2. However, as seen in the Example in Section 3, simple relationships between explanatory variables and cause-specific hazards do not lead to simple relationships between explanatory variables and cumulative incidences. Thus, though roughly identical relapse rates were seen for EBMT risk groups 0/1 and 2, the risk of relapse was higher for EBMT risk group 0/1 due to a lower rate of NRM for that group.

Such properties have motivated the development of models that directly link the cumulative incidence to explanatory variables. The most popular

model of this kind was introduced by Fine and Gray⁵ and links the cumulative incidence to explanatory variables in the same way as does the Cox model for all-cause mortality, see Box 4.

Example (continued)

Table 3 shows the result of the Fine-Gray regression model for relapse and NRM.

[Table 3 about here.]

The most striking aspect is the fact that the regression coefficient of EBMT risk group 2 for relapse is less than 0, albeit not statistically significant. This means that the cumulative incidence of relapse of EBMT risk group 2 is less than that of EBMT risk group 0/1. This is in agreement with Figure 5, although that was derived from a proportional cause-specific hazards model.

While the *relative sizes* of the regression coefficients in the Fine-Gray model in a useful way reflect the *ordering* of the cumulative incidence curves, their numerical values do not possess a simple interpretation. Thus, the estimates in Table 3 are *sub-distribution hazard ratios* and while this sounds like a hazard ratio, it is not. As noted above, the cause j -specific hazard gives the rate of cause j failure per time unit for individuals who are still alive. On the other hand, the cause j sub-distribution hazard gives the rate of cause j failure per time unit for individuals who are *either* still alive *or have already died from causes other than j* ⁵. Thus, a sub-distribution hazard bears no resemblance to an epidemiological rate, since individuals who have died from another cause remain in the risk set, even though they are no longer at risk of

experiencing a cause j failure. This fact does complicate the interpretation of parameters from the Fine-Gray model.

A final, technical note is that the structure assumed in a Cox model for the cause-specific hazards (“proportional hazards”) is incompatible with that of the Fine-Gray model (“proportional sub-distribution hazards”)⁷. This means that careful checking of the model assumptions is important, both when inference is based on Cox models for cause-specific hazards and when it is based on Fine-Gray models.

Model checking may, initially, be performed graphically. Figure 6 shows non-parametric estimates of the cumulative incidences of relapse and non-relapse mortality for each of the five EBMT risk groups. The estimates of the relapse cumulative incidence curves for EBMT risks groups 0/1 and 2 cross. In the same way as crossing survival curves for two groups in all-cause mortality are an indication that the proportional hazards assumption may be violated, this suggests that the proportionality assumption of the sub-distribution hazards for relapse in the Fine-Gray model may be violated for EBMT risks groups 0/1 and 2.

[Figure 6 about here.]

A similar graphical way of checking the proportional hazards assumption of the proportional cause-specific hazards model is obtained by inspecting the non-parametric cause-specific hazard estimates of Figure 4. Also here the proportional hazards assumption is questionable, although (also for the Fine-Gray model) it seems that only the EMBT risk group 0/1 for relapse is causing the non-proportionality. On the other hand, crossing of the estimated curves could just signal that the true functions are identical and the graphical

examination can be complemented by formal significance testing, e.g. via the scaled Schoenfeld residuals as in a standard Cox model⁶ or following the lines of Andersen and Pohar Perme¹.

5 "Independent" competing risks

Throughout, we have discussed the "rate of failure from cause j " or the "risk of failure from cause j " but never the "time to failure from cause j ". This is because for some individuals failure from cause j will never occur and thereby, formally, allowing "time to failure from cause j " to be infinite. In contrast, a classical approach to competing risks is via latent failure times as briefly summarized by Kalbfleisch and Prentice⁹ (see Section 8.2.4). In that approach one imagines the existence of random variables, L_1, L_2 , representing time to failure from cause 1 and time to failure from cause 2, respectively. The data then include the smaller of L_1 and L_2 ($T = \text{time to failure}$) and the cause of failure (1 if $T = L_1$, 2 if $T = L_2$). (Right-censoring may be accounted for.) While this, at a first glance, may seem an attractive idea, the problem is that the distribution of (L_1, L_2) cannot be identified from the data described in Box 3 without further, unverifiable, assumptions^{3;11}. The typical such assumption is "independent" competing risks, that is, independence between L_1 and L_2 . This means that our incomplete observations of, e.g. L_1 , are the same as those that would have been observed in a hypothetical population where cause 2 is not operating. Since such an assumption is completely unverifiable based on data from this world (where cause 2 is, indeed, operating), we believe that the concept of "independent" competing risks is quite elusive and that analyses relying on "independence" should be

interpreted with great care.

Finally, note that the likelihood factorization in Box 3 should not be taken to be relying on an assumption of "independence" between the competing causes of failure. This factorization (and the resulting consequence that rates of cause 1 may be analyzed by, formally, treating cause 2 events as censorings and vice versa) solely relies on the definition of cause-specific hazards as the time-local rates of occurrence of events that are mutually exclusive.

6 Discussion

In epidemiology, rates and risks are frequently used as measures of disease incidence and in this paper we have re-iterated the fact that, in studies of all-cause mortality, they are equivalent due to their one-to-one correspondence, see Box 1. However, while both concepts generalize quite simply to the competing risks situation (rates are now cause-specific hazards and risks are cumulative incidences), a one-to-one correspondence between a single rate and the corresponding risk no longer exists. Thus, any given cumulative incidence depends on all cause-specific hazards (see Box 2) and vice versa, and even though a single "sub-distribution hazard" may be derived from a single cumulative incidence, this is not a rate in any standard epidemiological sense. Similarly, a "risk-type quantity" may formally be defined by plugging in a cause-specific hazard into the relationship derived in Box 1. However, the resulting "risk" may only be interpreted in a completely hypothetical world where the competing risk does not exist, see Section 5. This is also illustrated by the fact that the Kaplan-Meier estimator provides a biased estimate of the cumulative incidence in the presence of competing risks as

demonstrated in our example.

Another consequence of the lack of a one-to-one correspondence between rate and risk in a competing risks setting is that covariates may affect the cause- j specific hazard and the cause- j cumulative incidence differently. This was also illustrated in our example showing that, when it comes to regression modeling, there is a choice to be made whether models should focus on cause-specific hazards or on cumulative incidences. Cox regression models for cause-specific hazards have the advantage that they are easy to fit (simply censor for competing events, see Box 3) and they provide parameter estimates which possess simple rate ratio interpretations. Such models, however, do not provide simple relationships between covariates and the easier interpretable cumulative incidences. Such simple relationships may be obtained from Fine-Gray models but the price to be paid is a set of parameter estimates which are harder to interpret, see Box 4.

These properties, together with assessment of model fit, should be kept in mind when deciding on how to make inference in a competing risks situation. We believe, as also illustrated by our example, that both rates and risks remain useful and tend to supplement each other when studying models for competing risks. Cause-specific hazards may be more relevant when the disease etiology is of interest, since it quantifies the event rate among the ones at risk of developing the event of interest. Cumulative incidences are easier to interpret and are more relevant for the purpose of prediction.

Acknowledgements

This work was supported by the National Cancer Institute [grant number R01-54706-12 to P.K.A.], the Danish Natural Science Research Council [grant number 272-06-0442 "Point process modelling and statistical inference" to P.K.A.], and by the Netherlands Organization for Scientific Research [grant number ZONMW-912-07-018 "Prognostic modeling and dynamic prediction for competing risks and multi-state models" to H.P.].

References

- [1] Andersen, P. K. and Pohar Perme, M. 2010, 'Pseudo-observations in survival analysis', *Statistical Methods in Medical Research* **19**, 71–99.
- [2] Cox, D. R. 1972, 'Regression models and life-tables', *Journal of the Royal Statistical Society B* **34**, 187–220.
- [3] Cox, D. R. 1959, 'The analysis of exponentially distributed life-times with two types of failure', *Journal of the Royal Statistical Society B* **21**, 411–421.
- [4] dos Santos Silva, I. 1999, *Cancer Epidemiology: Principles and Methods*, Lyon, France: International Agency for Research on Cancer.
- [5] Fine, J. P. and Gray, R. J. 1999, 'A proportional hazards model for the subdistribution of a competing risk', *Journal of the American Statistical Association* **94**, 496–509.
- [6] Geskus, R. B. 2011, 'Cause-specific cumulative incidence estimation and

the fine and gray model under both left truncation and right censoring', *Biometrics* **67**.

- [7] Grambauer, N., Schumacher, M. and Beyersmann, J. 2010, 'Proportional subdistribution hazards modeling offers a summary analysis, even if misspecified', *Statistics in Medicine* **29**, 875–884.
- [8] Gratwohl, A., Hermans, J., Goldman, J. M., W., A., Carreras, E., Divergie, A., Frassoni, F., Gahrton, G., Kolb, H. J., Niederwieser, D., Ruutu, T., Vernant, J. P., de Witte, T. and Apperley, J. 1998, 'Risk assessment for patients with chronic myeloid leukaemia before allogeneic blood or marrow transplantation', *Lancet* **352**, 1087–1092.
- [9] Kalbfleisch, J. D. and Prentice, R. L. 2002, *The Statistical Analysis of Failure Time Data (2nd ed.)*, New York: Wiley.
- [10] Rothman, K. J. 2002, *Epidemiology: An Introduction*, New York: Oxford University Press.
- [11] Tsiatis, A. A. 1975, 'A nonidentifiability aspect of the problem of competing risks', *Proc. Nat. Acad. Sci. USA* **72**, 20–22.

Consider censored data in an all-cause mortality study consisting of independent observations $(T_1, D_1), \dots, (T_n, D_n)$ on n individuals. Here, T_i is the time of observation for individual i : a failure time if $D_i = 1$ and a time of censoring if $D_i = 0$. If the hazard rate function for the distribution of the true failure time for i is $h_i(t)$ and the cumulative hazard is $H_i(t)$, then the likelihood under independent censoring is

$$L_a = h_1(T_1)^{D_1} e^{-H_1(T_1)} \dots h_n(T_n)^{D_n} e^{-H_n(T_n)},$$

or written using the product symbol \prod :

$$L_a = \prod_{i=1}^n h_i(T_i)^{D_i} e^{-H_i(T_i)}.$$

Consider now competing risks data $(T_1, D_1), \dots, (T_n, D_n)$ with T_i still denoting the possibly censored time to failure and $D_i = 1$ if i is observed to fail from cause 1, $D_i = 2$ if i is observed to fail from cause 2, and $D_i = 0$ if i is censored. If the (cumulative) cause-specific hazard for individual i and cause $j = 1, 2$ is $(H_{ij}(t))$ $h_{ij}(t)$ then the likelihood under independent censoring is

$$L_c = \prod_{i=1}^n h_{i1}(T_i)^{I(D_i=1)} h_{i2}(T_i)^{I(D_i=2)} e^{-H_{i1}(T_i) - H_{i2}(T_i)},$$

which can be re-written as

$$L_c = L_{a1} L_{a2},$$

where

$$L_{aj} = \prod_{i=1}^n h_{ij}(T_i)^{I(D_i=j)} e^{-H_{ij}(T_i)}, j = 1, 2.$$

It follows that the likelihood for cause 1 is the same as it would have been in an all-cause mortality study considering only cause 1 failures and censoring when individuals fail from cause 2.

Box 3: Likelihood factorization in a competing risks study.

The Cox regression model specifies the hazard rate for given explanatory variables Z as

$$h(t | Z) = h_0(t) \exp(\beta Z),$$

where the baseline hazard $h_0(t)$ is the hazard rate when $Z = 0$. For an all-cause mortality study the risk of failure before time t is

$$F(t) = 1 - \exp(-H(t))$$

(see Box 1). It follows that for a Cox model

$$-\log(1 - F(t | Z)) = H_0(t) \exp(\beta Z),$$

where $H_0(t)$ is the cumulative baseline hazard.

The Fine-Gray model specifies the same sort of dependence on explanatory variables for the cumulative cause j incidence in a competing risks model, i.e.

$$-\log(1 - F_j(t | Z)) = \widetilde{H}_0(t) \exp(B_j Z).$$

As explained in the text, the right-hand side of this equation is the cumulative cause j sub-distribution hazard and as a consequence, the regression coefficients $\exp(B_j)$ are sub-distribution hazard ratios.

Box 4: Relationship between covariates and risk in the Cox model for all-cause mortality and in the Fine-Gray model for competing risks.

List of Figures

1	Nelson-Aalen estimates of the cumulative hazards (left) and the Kaplan-Meier estimates of the survival curves (right) for RFS for each of the five EBMT risk groups.	25
2	Naive Kaplan-Meier estimates of relapse and non-relapse mortality, shown as incidence and survival curves, respectively. . .	26
3	Cumulative incidence estimates of relapse and non-relapse mortality, shown as incidence and survival curves, respectively; naive Kaplan-Meier estimates are shown in grey.	27
4	Nelson-Aalen estimates of the cumulative cause-specific hazards of relapse and non-relapse mortality for each of the five EBMT risk groups.	28
5	Model-based cumulative incidence estimates for relapse and non-relapse mortality for each of the five EBMT risk groups. .	29
6	Non-parametric cumulative incidence estimates for relapse and non-relapse mortality for each of the five EBMT risk groups. .	30




Figure 1: Nelson-Aalen estimates of the cumulative hazards (left) and the Kaplan-Meier estimates of the survival curves (right) for RFS for each of the five EBMT risk groups.




Figure 2: Naive Kaplan-Meier estimates of relapse and non-relapse mortality, shown as incidence and survival curves, respectively.




Figure 3: Cumulative incidence estimates of relapse and non-relapse mortality, shown as incidence and survival curves, respectively; naive Kaplan-Meier estimates are shown in grey.




Figure 4: Nelson-Aalen estimates of the cumulative cause-specific hazards of relapse and non-relapse mortality for each of the five EBMT risk groups.




Figure 5: Model-based cumulative incidence estimates for relapse and non-relapse mortality for each of the five EBMT risk groups.




Figure 6: Non-parametric cumulative incidence estimates for relapse and non-relapse mortality for each of the five EBMT risk groups.

List of Tables

1	Number of censored observations and number of events for relapse and non-relapse mortality (NRM) in each of the EBMT risk groups.	32
2	Cause-specific hazard ratios and 95% confidence intervals (CIs) of the EBMT risk groups for relapse and non-relapse mortality.	33
3	Estimated regression coefficients (B) and associated standard errors (SE) of the EBMT risk groups for relapse and non-relapse, for Fine-Gray regression.	34

EBMT risk group	Total	Relapse $n(\%)$	NRM $n(\%)$	Censored $n(\%)$
0,1	506	113 (22.3%)	94 (18.6%)	299 (59.1%)
2	1159	247 (21.3%)	323 (27.9%)	589 (50.8%)
3	1218	292 (24.0%)	404 (33.2%)	522 (42.9%)
4	745	193 (25.9%)	300 (40.3%)	252 (33.8%)
5,6,7	354	112 (31.6%)	169 (47.7%)	73 (20.6%)

Table 1: Number of censored observations and number of events for relapse and non-relapse mortality (NRM) in each of the EBMT risk groups.

EBMT	Relapse		NRM	
risk group	HR	95% CI	HR	95% CI
0, 1				
2	1.01	0.81–1.27	1.57	1.25–1.97
3	1.28	1.03–1.59	2.01	1.61–2.52
4	1.57	1.25–1.99	2.68	2.12–3.37
5, 6, 7	2.67	2.06–3.47	3.98	3.09–5.13

Table 2: Cause-specific hazard ratios and 95% confidence intervals (CIs) of the EBMT risk groups for relapse and non-relapse mortality.

EBMT risk group	Relapse		NRM	
	B	SE	B	SE
0, 1				
2	-0.068	0.111	0.443	0.116
3	0.072	0.108	0.661	0.114
4	0.161	0.117	0.906	0.118
5, 6, 7	0.439	0.135	1.185	0.131

Table 3: Estimated regression coefficients (B) and associated standard errors (SE) of the EBMT risk groups for relapse and non-relapse, for Fine-Gray regression.

Department of Biostatistics
University of Copenhagen
Øster Farimagsgade 5
P.O. Box 2099
1014 Copenhagen K
Denmark

- 10/1 Andersen, P.K. & Skrdal, A. "Biological" interaction from a statistical point of view.
- 10/2 Rosthøj, S., Keiding, N., Schmiegelow, K. Application of History-Adjusted Marginal Structural Models to Maintenance Therapy of Children with Acute Lymphoblastic Leukaemia.
- 10/3 Parner, E.T. & Andersen, P.K. Regression analysis of censored data using pseudo-observations.
- 10/4 Nielsen, T. & Kreiner, S. Course Evaluation and Development: What can Learning Styles Contribute?
- 10/5 Lange, T. & Hansen, J.V. Direct and Indirect Effects in a Survival Context.
- 10/6 Andersen, P.K. & Keiding, N. Interpretability and importance of functionals in competing risks and multi-state models.
- 10/7 Gerds, T.A., Kattan, M.W., Schumacher, M. & Yu, C. Estimating a time-dependent concordance index for survival prediction models with covariate dependent censoring.
- 10/8 Mogensen, U.B., Ishwaran, H. & Gerds, T.A. Evaluating random forests for survival analysis using prediction error curves.
- 11/1 Kreiner, S. Is the foundation under PISA solid? A critical look at the scaling model underlying international comparisons of student attainment.
- 11/2 Andersen, P.K., Geskus, R.B. & Putter, H. Competing risks in epidemiology: Possibilities and pitfalls.